

3. A cross section has width $w = 2\sqrt{x}$ and area

$$A(x) = s^2 = \left(\frac{w}{\sqrt{2}}\right)^2 = 2x. \text{ The volume is}$$

$$\int_0^4 2x \, dx = \left[x^2\right]_0^4 = 16.$$

4. A cross section has width $w = (2-x^2) - x^2 = 2-2x^2$ and

$$\text{area } A(x) = \pi r^2 = \pi \left(\frac{w}{2}\right)^2 = \pi(1-x^2)^2. \text{ The volume is}$$

$$\begin{aligned} \int_{-1}^1 (1-x^2)^2 \, dx &= \pi \int_{-1}^1 (x^4 - 2x^2 + 1) \, dx \\ &= \pi \left[\frac{1}{5}x^5 - \frac{2}{3}x^3 + x \right]_{-1}^1 = \frac{16}{15}\pi. \end{aligned}$$

5. The cross section has width $w = 2\sqrt{1-x^2}$ and area

$$A(x) = s^2 = w^2 = 4(1-x^2). \text{ The volume is}$$

$$\int_{-1}^1 4(1-x^2) \, dx = 4 \int_{-1}^1 (1-x^2) \, dx = 4 \left[x - \frac{1}{3}x^3 \right]_{-1}^1 = \frac{16}{3}.$$

6. A cross section has width $w = 2\sqrt{1-x^2}$ and area

$$A(x) = s^2 = \left(\frac{w}{\sqrt{2}}\right)^2 = 2(1-x^2). \text{ The volume is}$$

$$\int_{-1}^1 2(1-x^2) \, dx = 2 \int_{-1}^1 (1-x^2) \, dx = 2 \left[x - \frac{1}{3}x^3 \right]_{-1}^1 = \frac{8}{3}.$$

7. The solid is a right circular cone of radius 1 and height 2.

$$V = \frac{1}{3}Bh = \frac{1}{3}(\pi r^2)h = \frac{1}{3}(\pi \cdot 1^2) \cdot 2 = \frac{2}{3}\pi$$

Using integration: A cross section has radius $\left(1 - \frac{1}{2}x\right)$

and area $A(x) = \pi \left(1 - \frac{1}{2}x\right)^2$. The volume is

$$\begin{aligned} V &= \int_0^2 \pi \left(1 - \frac{1}{2}x\right)^2 \, dx = \pi \int_0^2 \left(\frac{x^2}{4} - x + 1\right) \, dx \\ &= \pi \left[\frac{x^3}{12} - \frac{x^2}{2} + x \right]_0^2 = \frac{2}{3}\pi. \end{aligned}$$

8. The solid is a right circular cone of radius 3 and height 2.

$$V = \frac{1}{3}Bh = \frac{1}{3}(\pi r^2)h = \frac{1}{3}(\pi \cdot 3^2) \cdot 2 = 6\pi$$

Using integration: A cross section has radius $\left(\frac{3y}{2}\right)$

and area $A(y) = \pi \left(\frac{3y}{2}\right)^2$. The volume is

$$\begin{aligned} V &= \int_0^2 \pi \left(\frac{3y}{2}\right)^2 \, dy = \frac{9}{4}\pi \int_0^2 y^2 \, dy \\ &= \frac{9}{4}\pi \left[\frac{y^3}{3} \right]_0^2 = 6\pi. \end{aligned}$$

9. A cross section has radius $r = \tan\left(\frac{\pi}{4}y\right)$ and area

$$A(y) = \pi r^2 = \pi \tan^2\left(\frac{\pi}{4}y\right). \text{ The volume is}$$

$$\begin{aligned} \int_0^1 \pi \tan^2\left(\frac{\pi}{4}y\right) \, dy &= \pi \left[\frac{4}{\pi} \tan\left(\frac{\pi}{4}y\right) - y \right]_0^1 \\ &= \pi \left(\frac{4}{\pi} - 1 \right) \\ &= 4 - \pi. \end{aligned}$$

10. A cross section has radius $r = \sin x \cos x$ and area


$$A(x) = \pi r^2 = \pi \sin^2 x \cos^2 x. \text{ The shaded region extends}$$

from $x = 0$ to where $\sin x \cos x$ drops back to 0, i.e., where

$$x = \frac{\pi}{2}. \text{ Now, since } \cos 2x = 2 \cos^2 x - 1, \text{ we know}$$

$$\cos^2 x = \frac{1 + \cos 2x}{2} \text{ and since } \cos 2x = 1 - 2 \sin^2 x, \text{ we}$$

$$\begin{aligned} \text{know } \sin^2 x &= \frac{1 - \cos 2x}{2}. \\ \int_0^{\pi/2} \pi \sin^2 x \cos^2 x \, dx &= \pi \int_0^{\pi/2} \frac{1 - \cos 2x}{2} \cdot \frac{1 + \cos 2x}{2} \, dx \\ &= \frac{\pi}{4} \int_0^{\pi/2} (1 - \cos^2 2x) \, dx = \frac{\pi}{4} \int_0^{\pi/2} \sin^2 2x \, dx \\ &= \frac{\pi}{4} \int_0^{\pi/2} \frac{1 - \cos 4x}{2} \, dx = \frac{\pi}{8} \int_0^{\pi/2} (1 - \cos 4x) \, dx \\ &= \frac{\pi}{8} \left[x - \frac{1}{4} \sin 4x \right]_0^{\pi/2} = \frac{\pi}{8} \left[\left(\frac{\pi}{2} - 0 \right) - 0 \right] = \frac{\pi^2}{16}. \end{aligned}$$




$[-2, 4]$ by $[-1, 5]$

A cross section has radius $r = x^2$ and area

$$A(x) = \pi r^2 = \pi x^4. \text{ The volume is}$$

$$\int_0^2 \pi x^4 \, dx = \pi \left[\frac{1}{5}x^5 \right]_0^2 = \frac{32\pi}{5}.$$




$[-4, 6]$ by $[-1, 9]$

A cross section has radius $r = x^3$ and area

$$A(x) = \pi r^2 = \pi x^6. \text{ The volume is}$$

$$\int_0^2 \pi x^6 \, dx = \pi \left[\frac{1}{7}x^7 \right]_0^2 = \frac{128\pi}{7}.$$



$[-6, 6]$ by $[-4, 4]$


The solid is a sphere of radius $r = 3$. The volume is

$$\frac{4}{3}\pi r^3 = 36\pi.$$

Using integration: A cross section has radius $\sqrt{9-x^2}$ and

area $A(y) = \pi(\sqrt{9-x^2})^2$. The volume is

$$\begin{aligned} V &= \int_{-3}^3 \pi(\sqrt{9-x^2})^2 dx = \pi \int_{-3}^3 (9-x^2) dx \\ &= \pi \left[9x - \frac{1}{3}x^3 \right]_{-3}^3 = 36\pi. \end{aligned}$$



$[-0.5, 1.5]$ by $[-0.5, 0.5]$

The parabola crosses the line $y = 0$ when


$x - x^2 = x(1-x) = 0$, i.e., when $x = 0$ or $x = 1$. A cross

section has radius $r = x - x^2$ and area

$$A(x) = \pi r^2 = \pi(x - x^2)^2 = \pi(x^2 - 2x^3 + x^4).$$

The volume is


$$\int_0^1 \pi(x^2 - 2x^3 + x^4) dx = \pi \left[\frac{1}{3}x^3 - \frac{1}{2}x^4 + \frac{1}{5}x^5 \right]_0^1 = \frac{\pi}{30}.$$



$[-1, 2]$ by $[-1, 2]$

Use cylindrical shells: A shell has radius y and height y .


$$\text{The volume is } \int_0^1 2\pi(y)(y) dy = 2\pi \left[\frac{1}{3}y^3 \right]_0^1 = \frac{2}{3}\pi.$$



$[-1, 3]$ by $[-1, 3]$

Use washer cross sections: A washer has inner radius $r = x$, outer radius $R = 2x$, and area $A(x) = \pi(R^2 - r^2) = 3\pi x^2$.

$$\text{The volume is } \int_0^1 3\pi x^2 dx = 3\pi \left[\frac{1}{3}x^3 \right]_0^1 = \pi.$$




$[-2, 3]$ by $[-1, 6]$

The curves intersect when $x^2 + 1 = x + 3$, which is when $x^2 - x - 2 = (x-2)(x+1) = 0$, i.e., when $x = -1$ or $x = 2$.

Use washer cross sections: a washer has inner radius $r = x^2 + 1$, outer radius $R = x + 3$, and area

$$\begin{aligned} A(x) &= \pi(R^2 - r^2) \\ &= \pi[(x+3)^2 - (x^2+1)^2] \\ &= \pi(-x^4 - x^2 + 6x + 8). \end{aligned} \text{ The volume is}$$

$$\begin{aligned} &\int_{-1}^2 \pi(-x^4 - x^2 + 6x + 8) dx \\ &= \pi \left[-\frac{1}{5}x^5 - \frac{1}{3}x^3 + 3x^2 + 8x \right]_{-1}^2 \\ &= \pi \left[\left(-\frac{32}{5} - \frac{8}{3} + 12 + 16 \right) - \left(\frac{1}{5} + \frac{1}{3} + 3 - 8 \right) \right] = \frac{117\pi}{5}. \end{aligned}$$



$[-2, 3]$ by $[-1, 5]$


The curves intersect when $4 - x^2 = 2 - x$, which is when $x^2 - x - 2 = (x-2)(x+1) = 0$, i.e., when $x = -1$ or $x = 2$.

Use washer cross sections: a washer has inner radius $r = 2 - x$, outer radius $R = 4 - x^2$, and area

$$\begin{aligned} A(x) &= \pi(R^2 - r^2) \\ &= \pi[(4-x^2)^2 - (2-x)^2] \\ &= \pi(12 + 4x - 9x^2 + x^4). \end{aligned}$$

The volume is

$$\begin{aligned} &\int_{-1}^2 \pi(12 + 4x - 9x^2 + x^4) dx \\ &= \pi \left[12x + 2x^2 - 3x^3 + \frac{1}{5}x^5 \right]_{-1}^2 \\ &= \pi \left[\left(24 + 8 - 24 + \frac{32}{5} \right) - \left(-12 + 2 + 3 - \frac{1}{5} \right) \right] = \frac{108\pi}{5}. \end{aligned}$$



$\left[-\frac{\pi}{3}, \frac{\pi}{3} \right]$ by $[-0.5, 2]$


Use washer cross sections: a washer has inner radius $r = \sec x$, outer radius $R = \sqrt{2}$, and area

$$A(x) = \pi(R^2 - r^2) = \pi(2 - \sec^2 x).$$

19. Continued

The volume is

$$\begin{aligned} \int_{-\pi/4}^{\pi/4} \pi(2 - \sec^2 x) dx &= \pi \left[2x - \tan x \right]_{-\pi/4}^{\pi/4} \\ &= \pi \left[\left(\frac{\pi}{2} - 1 \right) - \left(-\frac{\pi}{2} + 1 \right) \right] \\ &= \pi^2 - 2\pi. \end{aligned}$$




[-1, 5] by [-3, 1]

The curves intersect where $-\sqrt{x} = -2$, which is where $x=4$. Use washer cross sections: a washer has inner radius

$r = \sqrt{x}$, outer radius $R=2$, and area

$$A(x) = \pi(R^2 - r^2) = \pi(4 - x).$$

The volume is $\int_0^4 \pi(4 - x) dx = \pi \left[4x - \frac{1}{2}x^2 \right]_0^4 = 8\pi$




[-0.5, 1.5] by [-0.5, 2]

The curves intersect at $x = \frac{\pi}{4} \approx 0.7854$. A cross section has

radius $r = \sqrt{2} - \sec x \tan x$ and area

$A(x) = \pi r^2 = \pi(\sqrt{2} - \sec x \tan x)^2$. Use NINT to find

$$\int_0^{0.7854} \pi(\sqrt{2} - \sec x \tan x)^2 dx \approx 2.301.$$




[-1, 3] by [-1, 3]

The curve and horizontal line intersect at $x = \frac{\pi}{2}$. A cross section has radius $2 - 2 \sin x$ and area

$$A(x) = \pi r^2 = 4\pi(1 - \sin x)^2 = 4\pi(1 - 2 \sin x + \sin^2 x).$$

The volume is

$$\begin{aligned} \int_0^{\pi/2} 4\pi(1 - 2 \sin x + \sin^2 x) dx \\ &= \int_0^{\pi/2} 4\pi \left(1 - 2 \sin x + \frac{1 - \cos 2x}{2} \right) dx \\ &= 4\pi \left[\frac{3}{2}x + 2 \cos x - \frac{1}{4} \sin 2x \right]_0^{\pi/2} \\ &= 4\pi \left(\frac{3\pi}{4} - 2 \right) = \pi(3\pi - 8) \end{aligned}$$




[-1, 3] by [-1.5, 1.5]

A cross section has radius $r = \sqrt{5}y^2$ and area

$$A(y) = \pi r^2 = 5\pi y^4.$$

The volume is $\int_{-1}^1 5\pi y^4 dy = \pi \left[y^5 \right]_{-1}^1 = 2\pi$.




[-1, 4] by [-1, 3]

A cross section has radius $r = y^{3/2}$ and area

$$A(y) = \pi r^2 = \pi y^3. \text{ The volume is}$$

$$\int_0^2 \pi y^3 dy = \pi \left[\frac{1}{4} y^4 \right]_0^2 = 4\pi.$$




[-1.2, 3.5] by [-1, 2.1]

Use washer cross sections. A washer has inner radius $r = 1$. Outer radius $R = y + 1$, and area

$$A(y) = \pi(R^2 - r^2) = \pi[(y+1)^2 - 1] = \pi(y^2 + 2y). \text{ The}$$


volume is $\int_0^1 \pi(y^2 + 2y) dy = \pi \left[\frac{1}{3} y^3 + y^2 \right]_0^1 = \frac{4}{3} \pi$.



[-1.7, 3] by [-1, 2.1]

Use cylindrical shells: a shell has radius x and height x . The


$$\text{volume is } \int_0^1 2\pi(x)(x) dx = 2\pi \left[\frac{1}{3} x^3 \right]_0^1 = \frac{2}{3} \pi.$$



[-2, 4] by [-1, 5]

Use cylindrical shells: A shell has radius x and height x^2 .


$$\text{The volume is } \int_0^2 2\pi(x)(x^2) dx = 2\pi \left[\frac{1}{4} x^4 \right]_0^2 = 8\pi.$$



$[-0.5, 1.5]$ by $[-0.5, 1.5]$

The curves intersect at $x=0$ and $x=1$. Use cylindrical shells: a shell has radius x and height $\sqrt{x}-x$. The volume is

$$\int_0^1 2\pi(x)(\sqrt{x}-x)dx = 2\pi \left[\frac{2}{5}x^{5/2} - \frac{1}{3}x^3 \right]_0^1 = \frac{2\pi}{15}.$$



$[-1, 5]$ by $[-1, 3]$

The curved and horizontal line intersect at $(4, 2)$.

(a) Use washer cross sections: a washer has inner radius

$$r = \sqrt{x}, \text{ outer radius } R = 2, \text{ and area}$$

$$A(x) = \pi(R^2 - r^2) = \pi(4 - x). \text{ The volume is}$$

$$\int_0^4 \pi(4 - x)dx = \pi \left[4x - \frac{1}{2}x^2 \right]_0^4 = 8\pi$$

(b) A cross section has radius $r = y^2$ and area

$$A(y) = \pi r^2 = \pi y^4.$$

$$\text{The volume is } \int_0^2 \pi y^4 dy = \pi \left[\frac{1}{5}y^5 \right]_0^2 = \frac{32\pi}{5}.$$

(c) A cross section has radius $r = 2 - \sqrt{x}$ and area

$$A(x) = \pi r^2 = \pi(2 - \sqrt{x})^2 = \pi(4 - 4\sqrt{x} + x).$$

The volume is

$$\int_0^4 \pi(4 - 4\sqrt{x} + x)dx = \pi \left[4x - \frac{8}{3}x^{3/2} + \frac{1}{2}x^2 \right]_0^4 = \frac{8\pi}{3}.$$


(d) Use washer cross sections: a washer has inner radius

$$r = 4 - y^2, \text{ outer radius } R = 4, \text{ and area}$$

$$A(y) = \pi(R^2 - r^2) = \pi[16 - (4 - y^2)^2] = \pi(8y^2 - y^4).$$

The volume is

$$\int_0^2 \pi(8y^2 - y^4)dy = \pi \left[\frac{8}{3}y^3 - \frac{1}{5}y^5 \right]_0^2 = \frac{224\pi}{15}$$



$[-1, 3]$ by $[-1, 3]$

The slanted and vertical lines intersect at $(1, 2)$

(a) The solid is a right circular cone of radius 1 and height 2. The volume is

$$\frac{1}{3}Bh = \frac{1}{3}(\pi r^2)h = \frac{1}{3}(\pi 1^2)2 = \frac{2}{3}\pi.$$


Using integration: A cross section has radius $1 - \frac{y}{2}$

and area $A(y) = \pi r^2 = \pi \left(1 - \frac{y}{2}\right)^2$. The volume is

$$\begin{aligned} V &= \int_0^2 \pi \left(1 - \frac{y}{2}\right)^2 dy = \pi \int_0^2 \left(1 - y + \frac{1}{4}y^2\right) dy \\ &= \pi \left[y - \frac{1}{2}y^2 + \frac{1}{12}y^3 \right]_0^2 = \frac{2}{3}\pi. \end{aligned}$$

(b) Use cylindrical shells: shell has radius $2 - x$ and height $2x$. The volume is

$$\begin{aligned} \int_0^1 2\pi(2-x)(2x)dx &= 4\pi \int_0^1 (2x - x^2)dx \\ &= 4\pi \left[x^2 - \frac{1}{3}x^3 \right]_0^1 = \frac{8\pi}{3}. \end{aligned}$$



$[-2, 2]$ by $[-1, 2]$

The curves intersect at $(\pm 1, 1)$.

(a) A cross section has radius $r = 1 - x^2$ and area

$$A(x) = \pi r^2 = \pi(1 - x^2)^2 = \pi(1 - 2x^2 + x^4).$$

The volume is

$$\int_{-1}^1 \pi(1 - 2x^2 + x^4)dx = \pi \left[x - \frac{2}{3}x^3 + \frac{1}{5}x^5 \right]_{-1}^1 = \frac{16\pi}{15}.$$

(b) Use cylindrical shells: a shell has radius $2 - y$ and height $2\sqrt{y}$. The volume is

$$\begin{aligned} \int_0^1 2\pi(2-y)(2\sqrt{y})dy &= 4\pi \int_0^1 (2\sqrt{y} - y^{3/2})dy \\ &= 4\pi \left[\frac{4}{3}y^{3/2} - \frac{2}{5}y^{5/2} \right]_0^1 = \frac{56\pi}{15}. \end{aligned}$$

(c) Use cylindrical shells: a shell has radius $y+1$ and height $2\sqrt{y}$. The volume is

$$\begin{aligned} \int_0^1 2\pi(y+1)(2\sqrt{y})dy &= 4\pi \int_0^1 (y^{3/2} + \sqrt{y})dy \\ &= 4\pi \left[\frac{2}{5}y^{5/2} + \frac{2}{3}y^{3/2} \right]_0^1 = \frac{64\pi}{15}. \end{aligned}$$

32. (a) A cross section has radius $r = h\left(1 - \frac{x}{b}\right)$ and area

$$A(x) = \pi r^2 = \pi h^2 \left(1 - \frac{x}{b}\right)^2. \text{ The volume is}$$

$$\int_0^b \pi h^2 \left(1 - \frac{x}{b}\right)^2 dx = \pi h^2 \left[-\frac{b}{3} \left(1 - \frac{x}{b}\right)^3 \right]_0^b = \frac{\pi}{3} b h^2.$$

- (b) Use cylindrical shells: a shell has radius x and height

$$h\left(1 - \frac{x}{b}\right). \text{ The volume is}$$

$$\begin{aligned} \int_0^b 2\pi(x)h\left(1 - \frac{x}{b}\right) dx &= 2\pi h \int_0^b \left(x - \frac{x^2}{b}\right) dx \\ &= 2\pi h \left[\frac{1}{2}x^2 - \frac{x^3}{3b} \right]_0^b = \frac{\pi}{3} b^2 h. \end{aligned}$$

33. A shell has height $12(y^2 - y^3)$.

- (a) A shell has radius y . The volume is

$$\begin{aligned} \int_0^1 2\pi(y)12(y^2 - y^3) dy &= 24\pi \int_0^1 (y^3 - y^4) dy \\ &= 24\pi \left[\frac{1}{4}y^4 - \frac{1}{5}y^5 \right]_0^1 = \frac{6\pi}{5}. \end{aligned}$$

- (b) A shell has radius $1 - y$. The volume is

$$\begin{aligned} \int_0^1 2\pi(1 - y)12(y^2 - y^3) dy &= 24\pi \int_0^1 (y^4 - 2y^3 + y^2) dy \\ &= 24\pi \left[\frac{1}{5}y^5 - \frac{1}{2}y^4 + \frac{1}{3}y^3 \right]_0^1 = \frac{4\pi}{5}. \end{aligned}$$

- (c) A shell has radius $\frac{8}{5} - y$. The volume is

$$\begin{aligned} \int_0^1 2\pi\left(\frac{8}{5} - y\right)12(y^2 - y^3) dy &= 24\pi \int_0^1 \left(y^4 - \frac{13}{5}y^3 + \frac{8}{5}y^2\right) dy \\ &= 24\pi \left[\frac{1}{5}y^5 - \frac{13}{20}y^4 + \frac{8}{15}y^3 \right]_0^1 = 2\pi. \end{aligned}$$

- (d) A shell has radius $y + \frac{2}{5}$. The volume is

$$\begin{aligned} \int_0^1 2\pi\left(y + \frac{2}{5}\right)12(y^2 - y^3) dy &= 24\pi \int_0^1 \left(-y^4 + \frac{3}{5}y^3 + \frac{2}{5}y^2\right) dx \\ &= 24\pi \left[-\frac{1}{5}y^5 + \frac{3}{20}y^4 + \frac{2}{15}y^3 \right]_0^1 = 2\pi. \end{aligned}$$

34. A shell has height $\frac{y^2}{2} - \left(\frac{y^4}{4} - \frac{y^2}{2}\right) = y^2 - \frac{y^4}{4}$.

- (a) A shell has radius y . The volume is

$$\int_0^2 2\pi(y)\left(y^2 - \frac{y^4}{4}\right) dy = 2\pi \left[\frac{1}{4}y^4 - \frac{1}{24}y^6 \right]_0^2 = \frac{8\pi}{3}.$$

- (b) A shell has radius $2 - y$. The volume is


$$\begin{aligned} \int_0^2 2\pi(2 - y)\left(y^2 - \frac{y^4}{4}\right) dy &= 2\pi \int_0^2 \left(\frac{y^5}{4} - \frac{y^4}{2} - y^3 + 2y^2\right) dy \\ &= 2\pi \left[\frac{1}{24}y^6 - \frac{1}{10}y^5 - \frac{1}{4}y^4 + \frac{2}{3}y^3 \right]_0^2 = \frac{8\pi}{5}. \end{aligned}$$

- (c) A shell has radius $5 - y$. The volume is

$$\begin{aligned} \int_0^2 2\pi(5 - y)\left(y^2 - \frac{y^4}{4}\right) dy &= 2\pi \int_0^2 \left(\frac{y^5}{4} - \frac{5y^4}{4} - y^3 + 5y^2\right) dy \\ &= 2\pi \left[\frac{1}{24}y^6 - \frac{1}{4}y^5 - \frac{1}{4}y^4 + \frac{5}{3}y^3 \right]_0^2 = 8\pi. \end{aligned}$$

- (d) A shell has radius $y + \frac{5}{8}$. The volume is


$$\begin{aligned} \int_0^2 2\pi\left(y + \frac{5}{8}\right)\left(y^2 - \frac{y^4}{4}\right) dy &= 2\pi \int_0^2 \left(-\frac{y^5}{4} - \frac{5y^4}{32} + y^3 + \frac{5y^2}{8}\right) dy \\ &= 2\pi \left[-\frac{1}{24}y^6 - \frac{1}{32}y^5 + \frac{1}{4}y^4 + \frac{5}{24}y^3 \right]_0^2 = 4\pi. \end{aligned}$$



$[-2, 3]$ by $[-2, 3]$

A shell has radius x and height $x - \left(-\frac{x}{2}\right) = \frac{3}{2}x$.

The volume is $\int_0^2 2\pi(x)\left(\frac{3}{2}x\right) dx = \pi \left[x^3\right]_0^2 = 8\pi$.




$[-2, 2]$ by $[-1, 3]$

$x^2 = 2 - x$ at $x = 1$. A shell has radius x and height

$2 - x - x^2$. The volume is


$$\int_0^1 2\pi(x)(2 - x - x^2) dx = 2\pi \left[x^2 - \frac{1}{3}x^3 - \frac{1}{4}x^4 \right]_0^1 = \frac{5\pi}{6}.$$



$[-1, 5]$ by $[-1, 3]$

A shell has radius x and height \sqrt{x} . The volume is

$$\int_0^4 2\pi(x)(\sqrt{x}) dx = 2\pi \left[\frac{2}{5}x^{5/2} \right]_0^4 = \frac{128\pi}{5}.$$



$[-2, 2]$ by $[-2, 2]$

The functions intersect where $2x - 1 = \sqrt{x}$, i.e., at $x = 1$.

A shell has radius x and height

$\sqrt{x} - (2x - 1) = \sqrt{x} - 2x + 1$. The volume is

$$\begin{aligned} \int_0^1 2\pi(x)(\sqrt{x} - 2x + 1) dx &= 2\pi \int_0^1 (x^{3/2} - 2x^2 + x) dx \\ &= 2\pi \left[\frac{2}{5}x^{5/2} - \frac{2}{3}x^3 + \frac{1}{2}x^2 \right]_0^1 \\ &= \frac{7\pi}{15}. \end{aligned}$$

39. A cross section has width $w = 2\sqrt{\sin x}$.

(a) $A(x) = \frac{\sqrt{3}}{4}w^2 = \sqrt{3}\sin x$, and

$$\begin{aligned} V &= \int_0^\pi \sqrt{3}\sin x dx \\ &= \sqrt{3} \int_0^\pi \sin x dx \\ &= \sqrt{3}[-\cos x]_0^\pi \\ &= 2\sqrt{3}. \end{aligned}$$

(b) $A(x) = s^2 = w^2 = 4\sin x$, and

$$V = \int_0^\pi 4\sin x dx = 4 \int_0^\pi \sin x dx = 4[-\cos x]_0^\pi = 8.$$

40. A cross section has width $w = \sec x - \tan x$.

(a) $A(x) = \pi r^2 = \pi \left(\frac{w}{2} \right)^2 = \frac{\pi}{4}(\sec x - \tan x)^2$, and

$$\begin{aligned} V &= \int_{-\pi/3}^{\pi/3} \frac{\pi}{4}(\sec x - \tan x)^2 dx \\ &= \frac{\pi}{4} \int_{-\pi/3}^{\pi/3} (\sec^2 x - 2\sec x \tan x + \tan^2 x) dx \\ &= \frac{\pi}{4} [\tan x - 2\sec x + \tan x - x]_{-\pi/3}^{\pi/3} \\ &= \frac{\pi}{2} \left[\tan x - \sec x - \frac{1}{2}x \right]_{-\pi/3}^{\pi/3} \\ &= \frac{\pi}{2} \left[\left(\sqrt{3} - 2 - \frac{\pi}{6} \right) - \left(-\sqrt{3} - 2 + \frac{\pi}{6} \right) \right] \\ &= \pi\sqrt{3} - \frac{\pi^2}{6}. \end{aligned}$$

(b) $A(x) = s^2 = w^2 = (\sec x - \tan x)^2$, and

$$V = \int_{-\pi/3}^{\pi/3} (\sec x - \tan x)^2 dx, \text{ which by same method as}$$

in part (a) equals $4\sqrt{3} - \frac{2}{3}\pi$.

41. A cross section has width $w = \sqrt{5}y^2$ and area

$$\pi r^2 = \pi \left(\frac{w}{2} \right)^2 = \frac{5\pi}{4}y^4. \text{ The volume is}$$

$$\int_0^2 \frac{5\pi}{4}y^4 dy = \frac{\pi}{4} [y^5]_0^2 = 8\pi.$$

42. A cross section has width $w = 2\sqrt{1 - y^2}$ and area


$$\frac{1}{2}s^2 = \frac{1}{2}w^2 = 2(1 - y^2). \text{ This volume is}$$

$$\int_{-1}^1 2(1 - y^2) dy = 2 \left[y - \frac{1}{3}y^3 \right]_{-1}^1 = \frac{8}{3}.$$

43. Since the diameter of the circular base of the solid extends from $y = \frac{12}{2} = 6$ to $y = 12$, for a diameter of 6 and a radius of 3, the solid has the same cross sections as the right circular cone. The volumes are equal by Cavalieri's Theorem.

44. (a) The volume is the same as if the square had moved without twisting: $V = Ah = s^2h$.

(b) Still s^2h : the lateral distribution of the square cross sections doesn't affect the volume. That's Cavalieri's Volume Theorem.



$[-1, 3]$ by $[-1.4, 9.1]$

The functions intersect at $(2, 8)$.

(a) Use washer cross sections: a washer has inner radius


$$r = x^3, \text{ outer radius } R = 4x, \text{ and area}$$

$$A(x) = \pi(R^2 - r^2) = \pi(16x^2 - x^6). \text{ The volume is}$$

$$\int_0^2 \pi(16x^2 - x^6) dx = \pi \left[\frac{16}{3}x^3 - \frac{1}{7}x^7 \right]_0^2 = \frac{512\pi}{21}.$$

(b) Use cylindrical shells: a shell has a radius $8 - y$ and height $y^{1/3} - \frac{y}{4}$. The volume is

$$\begin{aligned} \int_0^8 2\pi(8-y) \left(y^{1/3} - \frac{y}{4} \right) dy \\ = 2\pi \int_0^8 \left(8y^{1/3} - 2y - y^{4/3} + \frac{y^2}{4} \right) dy \\ = 2\pi \left[6y^{4/3} - y^2 - \frac{3}{7}y^{7/3} + \frac{1}{12}y^3 \right]_0^8 = \frac{832\pi}{21}. \end{aligned}$$



$[-0.5, 1.5]$ by $[-0.5, 1.5]$

The functions intersect at $(0, 0)$ and $(1, 1)$.


(a) Use cylindrical shells: A shell has radius x and height

$$2x - x^2 - x = x - x^2. \text{ The volume is}$$

$$\int_0^1 2\pi(x)(x - x^2) dx = 2\pi \left[\frac{1}{3}x^3 - \frac{1}{4}x^4 \right]_0^1 = \frac{\pi}{6}.$$

(b) Use cylindrical shells: a shell has radius $1 - x$ and height $2x - x^2 - x = x - x^2$. The volume is

$$\begin{aligned} \int_0^1 2\pi(1-x)(x - x^2) dx &= 2\pi \int_0^1 (x^3 - 2x^2 + x) dx \\ &= 2\pi \left[\frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{1}{2}x^2 \right]_0^1 \\ &= \frac{\pi}{6}. \end{aligned}$$



$[-0.5, 2.5]$ by $[-0.5, 2.5]$

The intersection points are $\left(\frac{1}{4}, 1\right)$, $\left(\frac{1}{4}, 2\right)$, and $(1, 1)$.

(a) A washer has inner radius $r = \frac{1}{4}$, outer radius $R = \frac{1}{y^2}$,

$$\text{and area } \pi(R^2 - r^2) = \pi \left(\frac{1}{y^4} - \frac{1}{16} \right). \text{ The volume is}$$

$$\int_1^2 \pi \left(\frac{1}{y^4} - \frac{1}{16} \right) dy = \pi \left[-\frac{1}{3y^3} - \frac{1}{16}y \right]_1^2 = \frac{11\pi}{48}.$$

(b) A shell has radius x and height $\frac{1}{\sqrt{x}} - 1$. The volume is

$$\int_{1/4}^1 2\pi(x) \left(\frac{1}{\sqrt{x}} - 1 \right) dx = 2\pi \left[\frac{2}{3}x^{3/2} - \frac{1}{2}x^2 \right]_{1/4}^1 = \frac{11\pi}{48}.$$

48. (a) For $0 < x \leq \pi$, $x f(x) = \frac{x(\sin x)}{x} = \sin x$.

$$\text{For } x = 0, x f(x) = 0 \cdot 1 = \sin 0 = \sin x. \text{ so}$$

$$x f(x) = \sin x \text{ for } 0 \leq x \leq \pi.$$

(b) Use cylindrical shells: a shell has radius x and height y . The volume is $\int_0^\pi 2\pi xy dx$, which from

$$\text{part (a) is } \int_0^\pi 2\pi \sin x dx = 2\pi [-\cos x]_0^\pi = 4\pi.$$

49. (a) A cross section has radius $r = \frac{x}{12} \sqrt{36 - x^2}$ and area

$$A(x) = \pi r^2 = \frac{\pi}{144} (36x^2 - x^4). \text{ The volume is}$$

$$\begin{aligned} \int_0^6 \frac{\pi}{144} (36x^2 - x^4) dx &= \frac{\pi}{144} \left[12x^3 - \frac{1}{5}x^5 \right]_0^6 \\ &= \frac{36\pi}{5} \text{ cm}^3. \end{aligned}$$

$$(b) \left(\frac{36\pi}{5} \text{ cm}^3 \right) (8.5 \text{ g/cm}^3) \approx 192.3 \text{ g}.$$

50. (a) A cross section has radius $r = \sqrt{2y}$ and area

$$\pi r^2 = 2\pi y. \text{ The volume is } \int_0^5 2\pi y dy = \pi [y^2]_0^5 = 25\pi.$$

50. Continued

$$(b) V(h) = \int A(h) dh, \text{ so } \frac{dV}{dh} = A(h).$$


$$\therefore \frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt} = A(h) \cdot \frac{dh}{dt},$$

$$\text{so } \frac{dh}{dt} = \frac{1}{A(h)} \cdot \frac{dV}{dt}$$

$$\text{For } h = 4, \text{ the area is } 2\pi(4) = 8\pi,$$

$$\text{so } \frac{dh}{dt} = \frac{1}{8\pi} \cdot 3 \frac{\text{units}^3}{\text{sec}} = \frac{3}{8\pi} \frac{\text{units}^3}{\text{sec}}.$$

51. (a)



The remaining solid is that swept out by the shaded region in revolution. Use cylindrical shells: a shell has radius x and height $2\sqrt{r^2 - x^2}$. The volume is

$$\begin{aligned} & \int_{\sqrt{r^2-2}}^r 2\pi(x)(2\sqrt{r^2-x^2}) dx \\ &= 2\pi \left[-\frac{2}{3}(r^2-x^2)^{3/2} \right]_{\sqrt{r^2-4}}^r \\ &= -\frac{4}{3}\pi(-8) = \frac{32\pi}{3}. \end{aligned}$$

(b) The answer is independent of r .

52. Partition the appropriate interval in the axis of revolution and measure the radius $r(x)$ of the shadow region at these points. Then use an approximation such as the trapezoidal rule to estimate the integral $\int_a^b \pi r^2(x) dx$.

53. Solve $ax - x^2 = 0$: This is true at $x = a$. For revolution about the x -axis, a cross section has radius $r = ax - x^2$ and area

$$A(x) = \pi r^2 = \pi(ax - x^2)^2 = \pi(a^2x^2 - 2ax^3 + x^4).$$

The volume is

$$\begin{aligned} \int_0^a \pi(a^2x^2 - 2ax^3 + x^4) dx &= \pi \left[\frac{1}{3}a^2x^3 - \frac{1}{2}ax^4 + \frac{1}{5}x^5 \right]_0^a \\ &= \frac{1}{30}\pi a^5. \end{aligned}$$

For revolution about the y -axis, a cylindrical shell has radius x and height $ax - x^2$. The volume is

$$\int_0^a 2\pi(x)(ax - x^2) dx = 2\pi \left[\frac{1}{3}ax^3 - \frac{1}{4}x^4 \right]_0^a = \frac{1}{6}\pi a^4.$$

Setting the two volumes equal,

$$\frac{1}{30}\pi a^5 = \frac{1}{6}\pi a^4 \text{ yields } \frac{1}{30}a = \frac{1}{6}, \text{ so } a = 5.$$

54. The slant height Δs of a tiny horizontal slice can be written as $\Delta s = \sqrt{\Delta x^2 + \Delta y^2} = \sqrt{1 + (g'(y))^2} \Delta y$. So the surface area is approximated by the Riemann sum

$$\sum_{k=1}^n 2\pi g(y_k) \sqrt{1 + (g'(y))^2} \Delta y. \text{ The limit of that is the integral.}$$

$$55. g'(y) = \frac{dx}{dy} = \frac{1}{2\sqrt{y}}, \text{ and}$$

$$\begin{aligned} \int_0^2 2\pi \sqrt{y} \sqrt{1 + \left(\frac{1}{2\sqrt{y}}\right)^2} dy &= \int_0^2 \pi \sqrt{4y+1} dy \\ &= \left[\frac{\pi}{6}(4y+1)^{3/2} \right]_0^2 \\ &= \frac{13\pi}{3} \approx 13.614 \end{aligned}$$

$$56. g'(y) = \frac{dx}{dy} = y^2, \text{ and}$$

$$\begin{aligned} \int_0^1 2\pi \left(\frac{y^3}{3}\right) \sqrt{1+(y^2)^2} dy &= \frac{2}{3}\pi \left[\frac{1}{6}(1+y^4)^{3/2} \right]_0^1 \\ &= \frac{\pi}{9}(2\sqrt{2}-1) \approx 0.638. \end{aligned}$$

$$57. g'(y) = \frac{dx}{dy} = \frac{1}{2}y^{-1/2}, \text{ and}$$

$$\begin{aligned} \int_1^3 2\pi \left[y^{1/2} - \left(\frac{1}{3}\right)^{3/2} \right] \sqrt{1 + \left(\frac{1}{2}y^{-1/2}\right)^2} dy \\ = 2\pi \int_1^3 \left[y^{1/2} - \left(\frac{1}{3}\right)^{3/2} \right] \sqrt{1 + \frac{1}{4y}} dy. \end{aligned}$$

Using NINT, this evaluates to ≈ 16.110

$$58. g'(y) = \frac{dx}{dy} = \frac{1}{\sqrt{2y-1}}, \text{ and}$$

$$\begin{aligned} \int_{5/8}^1 2\pi \sqrt{2y-1} \sqrt{1 + \left(\frac{1}{\sqrt{2y-1}}\right)^2} dy \\ = 2\pi \int_{5/8}^1 \sqrt{2y} dy \\ = 2\sqrt{2}\pi \left[\frac{2}{3}y^{3/2} \right]_{5/8}^1 \\ = \frac{4\sqrt{2}}{3}\pi \left(1 - \frac{5}{16}\sqrt{\frac{5}{2}} \right) \approx 2.997. \end{aligned}$$

$$59. f'(x) = \frac{dy}{dx} = 2x, \text{ and}$$

$$\int_0^2 2\pi x^2 \sqrt{1+(2x)^2} dx = \int_0^2 2\pi x^2 \sqrt{1+4x^2} dx \text{ evaluates, using NINT, to } \approx 53.226.$$

60. $f'(x) = \frac{dy}{dx} = 3 - 2x$, and

$$\int_0^3 2\pi(3x - x^2)\sqrt{1 + (3x - 2x)^2} dx \text{ evaluates, using NINT,}$$

to ≈ 44.877 .

61. $f'(x) = \frac{dy}{dx} = \frac{1-x}{\sqrt{2x-x^2}}$, and

$$\begin{aligned} \int_{0.5}^{1.5} 2\pi\sqrt{2x-x^2} \sqrt{1 + \left(\frac{1-x}{\sqrt{2x-x^2}}\right)^2} dx &= 2\pi \int_{0.5}^{1.5} 1 dx \\ &= 2\pi [x]_{0.5}^{1.5} \\ &= 2\pi \approx 6.283 \end{aligned}$$

62. $f'(x) = \frac{dy}{dx} = \frac{1}{2\sqrt{x+1}}$, and

$$\begin{aligned} \int_1^5 2\pi\sqrt{x+1} \sqrt{1 + \left(\frac{1}{2\sqrt{x+1}}\right)^2} dx \\ &= 2\pi \int_1^5 \sqrt{x + \frac{5}{4}} dx \\ &= 2\pi \left[\frac{2}{3} \left(x + \frac{5}{4}\right)^{3/2} \right]_1^5 \\ &= \frac{4\pi}{3} \left[\left(\frac{25}{4}\right)^{3/2} - \left(\frac{9}{4}\right)^{3/2} \right] = \frac{49\pi}{3} \approx 51.313 \end{aligned}$$

63. True. by definition

64. False. The volume is given by $\int_0^2 \pi y^4 dy$.

65. A. $V = \int_1^e (\ln(x))^2 dx = 0.718$

66. E. $V = \int_0^4 (\pi(8 - x^{3/2})) dx = 361.9$

67. B. $V = \int_0^{16} \pi(4^2 - (\sqrt{y})^2) dy = 128\pi$

68. D.

69. A cross section has radius $r = |c - \sin x|$ and area

$$A(x) = \pi r^2 = \pi(c - \sin x)^2 = \pi(c^2 - 2c \sin x + \sin^2 x).$$

The volume is

$$\begin{aligned} \int_0^\pi \pi(c^2 - 2c \sin x + \sin^2 x) dx \\ &= \pi \left[c^2 x - 2c \cos x + \frac{1}{2} x - \frac{1}{4} \sin 2x \right]_0^\pi \\ &= \pi \left[\left(c^2 \pi - 2c + \frac{1}{2} \pi \right) - 2c \right] \\ &= \pi^2 c^2 - 4\pi c + \frac{\pi^2}{2}. \end{aligned}$$

(a) Solve

$$\begin{aligned} \frac{d}{dc} \left[\pi^2 c^2 - 4\pi c + \frac{\pi^2}{2} \right] &= 0 \\ 2\pi^2 c - 4\pi &= 0 \\ \pi c - 2 &= 0 \\ c &= \frac{2}{\pi} \end{aligned}$$

This value of c gives a minimum for V because

$$\frac{d^2 V}{dc^2} = 2\pi^2 > 0.$$

Then the volume is $\pi^2 \left(\frac{2}{\pi}\right)^2 - 4\pi \left(\frac{2}{\pi}\right) + \frac{\pi^2}{2} = \frac{\pi^2}{2} - 4$


(b) Since the derivative with respect to c is not zero

anywhere else besides $c = \frac{2}{\pi}$, the maximum must occur

at $c = 0$ or $c = 1$. The volume for $c = 0$ is $\frac{\pi^2}{2} \approx 4.935$,

and for $c = 1$ it is $\frac{\pi(3\pi - 8)}{2} \approx 2.238$. $c = 0$ maximizes

the volume.



$[0, 2]$ by $[0, 6]$

The volume gets large without limit. This makes sense, since the curve is sweeping out space in an ever-increasing radius.

70. (a) Using $d = \frac{C}{\pi}$, and $A = \pi \left(\frac{d}{2}\right)^2 = \frac{C^2}{4\pi}$ yields the

following areas (in square inches, rounded to the nearest tenth): 2.3, 1.6, 1.5, 2.1, 3.2, 4.8, 7.0, 9.3, 10.7, 10.7, 9.3, 6.4, 3.2.

(b) If $C(y)$ is the circumference as a function of y , then the area of a cross section is

$$A(y) = \pi \left(\frac{C(y)/\pi}{2} \right)^2 = \frac{(C(y))^2}{4\pi}, \text{ and the volume is}$$

$$\frac{1}{4\pi} \int_0^6 (C(y))^2 dy.$$

$$\begin{aligned} \text{(c) } \frac{1}{4\pi} \int_0^6 A(y) dy &= \frac{1}{4\pi} \int_0^6 (C(y))^2 dy \\ &\approx \frac{1}{4\pi} \left(\frac{6-0}{24} \right) [5.4^2 + 2(4.5^2 + 4.4^2 \\ &\quad + 5.1^2 + 6.3^2 + 7.8^2 + 9.4^2 + 10.8^2 + 11.6^2 \\ &\quad + 11.6^2 + 10.8^2 + 9.0^2) + 6.3^2] \approx 34.7 \text{ in.}^3 \end{aligned}$$